

Tree Species Composition Influences Soil CO₂ and N₂O Fluxes in Amazon Forest

Joost van Haren^{1,2}, Scott Saleska², Alfredo Huete¹, Michael Keller³, Raimundo Cosme de Oliveira⁴


¹Soil Water and Environmental Science, University of Arizona

²Ecology and Evolutionary Biology, University of Arizona

³University of New Hampshire/US Forest Service, Puerto Rico

⁴Embrapa Occidental, Belem

Why measure tropical soil CO₂ and N₂O fluxes?



Contribution of
tropical soils to
global budget

CO₂: largest soil
source

(Raich and
Schlesinger, 1992)

N₂O: ~20% all natural
sources


(Bouwman et al.,
1993)

Biogeochemical models

- Include:
 - Temperature
 - NGAS (Parton et al. 1996)
 - Precipitation
 - ECOSYS (Grant & Pattey, 1999)
 - Soil Parameters
 - Century (Liu et al. 2000)
 - PnET-N-DCDN (Li et al. 2000)
 - NASA-CASA (Potter et al. 2001)
 - Moisture
 - Texture
 - C/N
 - Substrate quality and quantity

Tree species are not explicitly considered


Plants influence on soil C



and available C on heterotrophic denitrifiers

Garcia-Montiel et al. (2003)
added water and glucose to forest soil in Rondonia

Glucose amplifies
N₂O fluxes



Hypotheses




- **Plant species traits and biomass significantly influence soil CO₂ and N₂O fluxes.**
- **Satellite based remote sensing could help scale trace gas fluxes, through detecting species traits (such as leaf N content) and biomass.**

Tapajós National Forest




Belterra Plantation

- Mature trees planted in 1978
- Pure and mixed species plots
- Soil texture: >80% clay and < 5% sand



Km 67 Tower location

- Standing Biomass inventory since 1999
- Very diverse: 263 tree species
- Soil texture: >80% clay and < 5% sand




Field measurements

- 167 flux locations near 14 tree species
- Gas flux
- air and soil T
- 0-3 cm BD and % WFPS
- DBH of main tree and DBH and species of all trees > 1 cm within a 3 m radius





Tree species (C = canopy, E = emergent, P = pioneer)		Base area % of total
<i>Caryocar villosum</i> (Piquiá, E)	PI	3.3
<i>Bertholletia excelsa</i> (Castanha de para, E)	CP	2.1
<i>Chamaecrista</i> sp. xinguensis (Coração de negro, C)	CN	6.2
<i>Couratari stellata</i> (Tauari, E)	TA	6.4
<i>Erismia uncinatum</i> (Quarubarana, E)	QU	16.3
<i>Lecythis lurida</i> (Jarana, E)	JA	7.0
<i>Manilkara huberi</i> (Maçaranduba, E)	MA	9.6
<i>Schefflera morototoni</i> (Morototo, P)	MO	0.5
<i>Pseudopiptadenia psilostachya</i> (Fava folha fina, E)	FF	2.3
<i>Sclerolobium chrysophyllum</i> (Tachi vermelho, C)	TV	3.5
<i>Pouteria reticulata</i> (Abiu, C)	AB	1.4
<i>Vochysia maxima</i> (Quaruba verdadeira, E)	QV	3.6
<i>Astronium lecointei</i> (Aroeira, C)	AR	0.74
<i>Copaifera multijuga</i> (Copaíba, C)	CO	3.0

Km 67 Forest Fluxes





Data do not support that forest tree species composition alters soil CO₂ flux



- Soil N₂O fluxes are different from overall mean
- Soil N₂O flux close to PI > QV, QU, and CO (at $\alpha = 0.05$ using Tukey-Kramer HSD test)

Tree species

Effects of biological and physical parameters




Linear relationship

Intercept = 2.45


Slope = 0.036

adjusted r² = 0.25



Soil N_2O flux close to
 Pi > Qv, Qu, and Co
 (at $\alpha = 0.05$ using Tukey-
 Kramer HSD test)

Correct for %WFPS



Soil N_2O flux close to
 Pi and Ab > Qu
 (at $\alpha = 0.05$ using Tukey-
 Kramer HSD test)

 Mean \pm 95% CI

Pi Ja Ab Cn Ar Cp Ta Tv Ma FF Mo Qv Qu Co
Tree species

Forest conclusions and last note

1. Soil CO₂ emissions appear to be unaffected by tree species composition
2. Soil N₂O emissions are consistently higher close to *Caryocar villosum* (Piquiá) relative to *Erisma uncinatum* (Quarubarana).
3. Even when corrected for %WFPS




N ₂ O flux μg-N m ⁻² hr ⁻¹	Mean	-95%	+95%
Grand mean	71.9	63.5	81.3
Species means	67.9	45.4	102.1
Background mean	38.9	50.4	30.0

Forest conclusions and last note

1. Soil CO₂ emissions appear to be unaffected by tree species composition
2. Soil N₂O emissions are consistently higher close to *Caryocar villosum* (Piquiá) relative to *Erisma uncinatum* (Quarubarana).
3. Even when corrected for %WFPS


N ₂ O flux μg-N m ⁻² hr ⁻¹	Mean	-95%	+95%
Grand mean	71.9	63.5	81.3
Species means	67.9	45.4	102.1
Background mean	38.9	50.4	30.0

However:





The relationship between %WFPS and ln(N₂O) might be species dependent

Belterra Plantation Fluxes





Soil CO₂ fluxes under
MA and PI > CP and
JA



Soil N₂O fluxes under
Pi > JA and AM




Effects of physical parameters



No correlation between
%WFPS and ln(N₂O)


Remote sensing

Leaf reflectance of key tree species in the Tapajos Forest are different at known N and protein absorption wavelengths



Leaf samples collected and measured with handheld ASD hyperspectral spectrometer by Timoaki Miura


Mean leaf reflectance spectra



Arrows indicate key N and protein Absorption bands

Remote sensing

Leaf reflectance of key tree species in the Tapajos forest are different at known N and protein absorption wavelengths




Leaf samples collected and measured with handheld ASD hyperspectral spectrometer by Timoaki Miura



+

Tree species means

Using PLS regression with spectral areas of known N and protein absorption features




Leaf samples collected and measured with handheld ASD hyperspectral spectrometer by Timoaki Miura


Tree species means

Using PLS regression with spectral areas of known N and protein absorption features

Remote sensing

Leaf reflectance of key tree species in the Tapajos forest are different at known N and protein absorption wavelengths





Conclusions

- Forest soil N_2O fluxes close to Piquia trees are greater than Quarubarana trees
- On the plantation tree species appear to influence both soil CO_2 and N_2O fluxes
- Hyperspectral techniques are able to distinguish tree species and could be used to predict plantation soil fluxes and determine soil N_2O 'hotspots' in forests.

This work would have been impossible without the fantastic help from the Santarem Field staff. Especially:

