

Carbon and Oxygen Isotope Ratio CO₂ Flux Analyses at the Soil, Canopy, and Landscape Scales

Co-PIs: Jim Ehleringer (Utah), Joe Berry (Carnegie), Luiz Martinelli (CENA-USP), Jean Ometto (CENA-USP)

collaborators: Antonio Nobre (INPA), Paulo Artaxo (USP)

CD-02

Abstract

We propose two areas of research emphasis to better understand and constrain estimates of carbon dynamics in Amazonia ecosystems, based on progress during Phase I of LBA-ECO.

(1) Stable isotope analyses of atmospheric CO₂ are one tool which scales processes from the soil and individual plant levels through to the troposphere. We will apply ¹³C/¹⁸O analyses as tools for better understanding gas exchange processes within ecosystems and regionally across ecosystems (forest, pasture, and river) in Manaus and Santarém. These data will also directly contribute to regional and aircraft studies which rely on ground observations for their data interpretation.

(2) Mechanistic leaf, canopy and site-scale canopy-scale models including isotope exchange are tools for extrapolating from local scale measurements towards understanding carbon cycling on an ecosystem and a basin-scale. Parameterized with leaf gas exchange characteristics (functional response curves, N values, ¹³C/¹⁸O and canopy structure) and driven by meteorological inputs, these models will be used to understand how climate influences carbon gain at the canopy-ecosystem scale and will be used to link with eddy covariance studies (NEE estimates), atmospheric sampling programs and regional carbon cycle modeling efforts as part of our science integration.

- The proposed study will make several key LBA-Ecology measurements at two primary regions: Manaus and Santarém:
- leaf-scale parameters to parameterize canopy photosynthesis models
 - canopy scale photosynthesis models driven by climate data
 - δ¹³C and δ¹⁸O values of stocks in forest and pastures: leaves and soils
 - δ¹³C and δ¹⁸O values of CO₂ effluxing from soils and from the total ecosystem at forest and pasture sites
 - δ¹³C and δ¹⁸O values of ecosystem photosynthetic discrimination in forests and pastures
 - δ¹³C and δ¹⁸O values of landscape-level respiration (PBL)

Our studies will be conducted in close association with eddy covariance measurements, atmospheric CO₂ sampling, and regional carbon modeling groups. The leaf and canopy-level models of carbon gain and of isotope fractionation will linked directly with regional scale models. The information gathered using our isotope studies will help in determining the role of the Amazon Basin as a net sink or source of CO₂ to the atmosphere.


Scientific objectives

- to use carbon and oxygen isotopes of both atmospheric CO₂ and organic matter to provide integrated information about important plant physiological characteristics across both spatial and temporal scales
- to use isotopes as a tool for understanding the mechanistic factors contributing to changes in NEE in response to climate drivers
- at the landscape scale, to use C and O isotopes to partition the relative contributions of the forest, pasture, and river components to CO₂ fluxes
- to couple mechanistic gas exchange data with models to predict the changes in stand-level photosynthesis associated with seasonal dynamics and with changes in precipitation and humidity

Canopy modeling studies

What have we done?

Quantifying photosynthetic parameters




What are we planning to do next?

Leaf and canopy models


The initial focus of our modeling work will be at the leaf scale. Our approach of building up from leaves to a canopy will test scaling assumptions and should lead to improved canopy models for larger scale simulations. The models we will use are derived from the leaf models of Collatz et al. (1991, 1992) extended to simulate isotopic fractionation in transpiration and photosynthesis. We will use within canopy meteorology obtained from the flux sites to drive these models at various levels within the forest or pasture canopies. We propose to collaborate with Scott Denning's group. They have developed a coupled model of physiology and turbulent transport within canopies, which provides a "bridge" to move from leaf-scale simulations to whole canopy simulations. The model will simulate within canopy gradients in CO₂ and its isotopic composition and can be tested against observations made in the field. Simulations with this multilayered model will be compared with simplified canopy models like SIB. We plan to use a new soil model developed by Riley et al. (2002) to simulate CO₂ and isotopic exchange associated with soil respiration and invasion of atmospheric CO₂ into the soil.

Isotope field studies

What have we done?




Isotopic composition of ecosystem respiration responds to environmental conditions, such as precipitation amount



Isotopic composition of daytime CO₂ reflects influence of both leaf (canopy) and soil sources


What are we planning to do next?

links to NEE through continuous observations



Organic C and O isotopes show limited seasonal dynamics, but there are pronounced profiles associated with stand structure

regional studies - forest, pasture, and river



regional studies - boundary layer

- balloon sampler to measure in PBL
- ¹³C and ¹⁸O of CO₂ in PBL profile
- PBL keeling plots
- ¹³C and ¹⁸O of CO₂ in forests
- ecosystem δ¹³C and δ¹⁸O discrimination



CD-02

Jim Ehleringer
ehleringer@biology.utah.edu

Luiz Martinelli
zebu@cena.usp.br

Joe Berry
joeberry@globalecology.stanford.edu

Jean Ometto
ometto@biology.utah.edu

Tomas Domingues
domingues@biology.utah.edu

Paulo Artaxo
artaxo@if.usp.br

Antonio Nobre
anobre27@yahoo.com